Wisconsin AIDS/HIV Surveillance Summary Cases reported 1983 through 2006

MILWAUKEE COUNTY

| | | ulative <u>Percent</u> | - | 2001-2005 Percent | | d in 2006 <u>Percent</u> | | ed alive <u>Percen</u> |
|---|---------------|---------------------------|--------|----------------------|-----------|-----------------------------|--------|---------------------------|
| Total cases | 4,789 | 100.0% | 884 | 100.0% | 183 | 100.0% | 3,010 | 100.0% |
| Current disease category | | | | | | | | |
| AIDS | 3,178 | 66.4% | 406 | 45.9% | 69 | 37.7% | 1,533 | 50.9% |
| Non-AIDS | 1,611 | 33.6% | 478 | 54.1% | 114 | 62.3% | 1,477 | 49.1% |
| Risk Exposure Categories | | | | | | | | |
| Men who have sex with men (MSM) | 2,472 | 51.6% | 387 | 43.8% | 84 | 45.9% | 1,423 | 47.39 |
| Injecting drug use (IDU) | 758 | 15.8% | 96 | 10.9% | 13 | 7.1% | 434 | 14.49 |
| MSM and IDU | 319 | 6.7% | 46 | 5.2% | 8 | 4.4% | 186 | 6.29 |
| Hemophilia/Coagulation disorder | 22 | 0.5% | 2 | 0.2% | 0 | 0.0% | 4 | 0.19 |
| High-risk heterosexual contact | 630 | 13.2% | 152 | 17.2% | 18 | 9.8% | 472 | 15.79 |
| Transfusion-associated | 18 | 0.4% | 4 | 0.5% | 0 | 0.0% | 9 | 0.39 |
| Mother with/at risk | 39 | 0.8% | 8 | 0.9% | 0 | 0.0% | 27 | 0.99 |
| Undetermined | 531 | 11.1% | 189 | 21.4% | 60 | 32.8% | 455 | 15.19 |
| Gender | | | | | | | | |
| Female | 863 | 18.0% | 220 | 24.9% | 30 | 16.4% | 645 | 21.49 |
| Male | 3,926 | 82.0% | 664 | 75.1% | 153 | 83.6% | 2,365 | 78.69 |
| Unknown | 0 | 0.0% | 0 | 0.0% | 0 | 0.0% | 0 | 0.0% |
| Dogo/Ethnicity | | | | | | | | |
| Race/Ethnicity | 4.070 | 20.20/ | 004 | 00.50/ | <i>EE</i> | 20.40/ | 000 | 20.00 |
| White | 1,876 | 39.2% | 261 | 29.5% | 55 | 30.1% | 980 | 32.69 |
| Black | 2,307 | 48.2% | 472 | 53.4% | 98 | 53.6% | 1,598 | 53.19 |
| Hispanic | 567 | 11.8% | 140 | 15.8% | 25 | 13.7% 1.6% | 405 | 13.5% |
| Asian/Pacific Islander | 13 | 0.3% 0.4% | 5 | 0.6% | 3 | 0.0% | 11 | 0.49 |
| American Indian | 19 | 0.4% 0.1% | 2 | 0.2% 0.5% | 0 | 0.0% | 9 | 0.39 |
| Multi-racial Unknown | 5 2 | 0.1% | 4 0 | 0.5% | 1 | 0.5% | 5 2 | 0.29 0.19 |
| | - | 0.070 | · · | 0.070 | · | 0.070 | _ | 0.17 |
| Age at diagnosis* | | | | | | | | |
| Under 5 | 37 | 0.8% | 7 | 0.8% | 1 | 0.5% | 3 | 0.19 |
| 5-14 | 13 | 0.3% | 2 | 0.2% | 0 | 0.0% | 15 | 0.5% |
| 15-19 | 135 | 2.8% | 41 | 4.6% | 9 | 4.9% | 15 | 0.5% |
| 20-24 | 531 | 11.1% | 103 | 11.7% | 31 | 16.9% | 87 | 2.9% |
| 25-44 | 3,463 | 72.3% | 584 | 66.1% | 106 | 57.9% | 1,571 | 52.29 |
| 45 years and older | 608 | 12.7% | 147 | 16.6% | 36 | 19.7% | 1,312 | 43.69 |
| Unknown | 2 | 0.0% | 0 | 0.0% | 0 | 0.0% | 4 | 0.19 |
| * For cases presumed alive, age is the age of | on 12/31/2006 | | | | | | | |
| Year of Report | | | | | | | | |
| Before 1990 | 726 | 15.2% | | | | | | |
| 1990 | 367 | 7.7% | | | | | | |
| 1991 | 349 | 7.3% | | | | | | |
| 1992 | 399 | 8.3% | | | | | | |
| 1993 | 326 | 6.8% | | | | | | |
| 1994 | 273 | 5.7% | | | | | | 1 - |
| 1995 | 272 | 5.7% | | | | | | |
| 1996 | 206 | 4.3% | | | | | | |
| 1997 | 234 | 4.9% | | | | | | |
| 1998 | 187 | 3.9% | | | | | | |
| 1999 | 186 | 3.9% | | | | | | |
| 2000 | 197 | 4.1% | | | | | | 7 |
| 2001 | 167 | 3.5% | | | | | | 7 |
| 2002 | 195 | 4.1% | | | | | | ı |
| 2002 | 158 | 3.3% | | | | | | |
| 2003 | 186 | | | | Ì | | | I. |
| | | 3.9% | | | | | | |
| 2005 | 178 | 3.7% | | | | | | |

183

3.8%

2006